

Metoclopramide Hydrochloride Oral Solution

5mg/5mL

Instructions for administration via nasogastric (NG) or percutaneous endoscopic gastrostomy (PEG) tubes:

Ensure that the enteral feeding tube is free from obstruction before administration.

1. Flush the enteral tube with 5mL of water.

2. Administer the required dose of Metoclopramide Hydrochloride Oral Solution with a suitable measuring device.

3. Flush the enteral tube with 5mL of water.

This product has not been tested with latex NG or PEG tubes and therefore should not be used with tubes made from latex.

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

Rosemont® 

DTM061.1 July 2020

Rosemont Pharmaceuticals Ltd. Rosemont House, Yorkdale Industrial Park, Braithwaite Street, Leeds LS11 9XE T +44 (0)113 244 1400 F +44 (0)113 245 3567 E infodesk@rosemontpharma.com Sales/ Customer Service: T +44 (0) 113 244 1999 F +44 (0) 113 246 0738 W www.rosemontpharma.com

Adverse events should be reported. Reporting forms and information can be found at www.mhra.gov.uk/yellowcard Adverse events should also be reported to Rosemont Pharmaceuticals Ltd on 0113 244 1400.