

Amlodipine Oral Solutions

1 mg/mL and 2 mg/mL

Instructions for administration via nasogastric (NG) or percutaneous endoscopic gastrostomy (PEG) tubes:

Amlodipine Oral Solution is suitable for use with the following types of NG and PEG tubes:

Material	External Bore Size (Fr Unit)	Internal Diameter (mm)	Maximum Length (cm)
Silicone	8	1.5	125
	10	2.0	125
PVC	8	1.5	125
	12	2.5	125
Polyurethane	8	1.5	125
	12	2.6	125
	18	4.0	75

Care should be taken during administration due to the viscous nature of the product.

Ensure that the enteral feeding tube is free from obstruction before administration.

The minimum volume of water that is required to flush tube sizes 8 Fr is 10mL and for tube sizes 10 Fr and above it is 20mL.

1. Flush the enteral tube with water, using the minimum flush volume required.

2. Administer the required dose of Amlodipine Oral Solution with a suitable measuring device.

3. Flush the enteral tube with water again using the minimum flush volume required.

This product has not been tested with latex NG or PEG tubes and therefore should not be used with tubes made from latex.

Air flushing of NG or PEG tubes to deliver the required dose of Amlodipine is not recommended.

The oral syringe provided in the pack should not be used for administration via NG or PEG tubes, HCPs must use another suitable device.

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.



DTM056.1 July 2020

Rosemont Pharmaceuticals Ltd. Rosemont House, Yorkdale Industrial Park, Braithwaite Street, Leeds LS11 9XE **T** +44 (0)113 244 1400 **F** +44 (0)113 245 3567 **E** infodesk@rosemontpharma.com **Sales/ Customer Service:** **T** +44 (0) 113 244 1999 **F** +44 (0) 113 246 0738 **W** www.rosemontpharma.com

Adverse events should be reported. Reporting forms and information can be found at www.mhra.gov.uk/yellowcard Adverse events should also be reported to Rosemont Pharmaceuticals Ltd on 0113 244 1400.