

Glycopyrronium Bromide Oral Solution

1 mg/5ml

Instruction for administration via nasogastric (NG) or percutaneous endoscopic gastrostomy (PEG) tubes:

Ensure that the enteral feeding tube is free from obstruction before administration.

1. Flush the enteral tube with water, using the minimum flush volume required.

2. Administer the required dose of glycopyrronium bromide oral solution with a suitable measuring device.

3. Flush the enteral tube again, using the minimum flush volume required.

Tubing size (French Units)	Recommended Minimum Flush Volume (ml)
4	1.2
6	2
8	5
10	8
12	10
18	10

With respect to tubal administration, this product should be administered with silicone, PVC, polyurethane NG or PEG tubes **only**.

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.



DTM324 October 2021

Rosemont Pharmaceuticals Ltd. Rosemont House, Yorkdale Industrial Park, Braithwaite Street, Leeds LS11 9XE **T** +44 (0)113 244 1400 **F** +44 (0)113 245 3567 **E** infodesk@rosemontpharma.com Sales/ Customer Service: **T** +44 (0) 113 244 1999 **F** +44 (0) 113 246 0738 **W** www.rosemontpharma.com

Adverse events should be reported. Reporting forms and information can be found at www.mhra.gov.uk/yellowcard Adverse events should also be reported to Rosemont Pharmaceuticals Ltd on 0113 244 1400.