

Omeprazole Powder for Oral Suspension

1mg/ml, 2mg/ml and 4mg/ml

Instructions for administration via nasogastric (NG) or percutaneous endoscopic gastrostomy (PEG) tubes:

Ensure that the enteral feeding tube is free from obstruction before administration.

1. Flush the enteral tube with 5mL of water.

2. Administer the required dose of Omeprazole Oral Suspension with a suitable measuring device.

3. Flush the enteral tube with 5mL of water.

This product is compatible for use with Polyurethane and PVC nasogastric (NG) and percutaneous endoscopic gastrostomy (PEG) tubes of size 6Fr to 16Fr.

1mg/ml & 2mg/ml Omeprazole Powder for Oral Suspension – for the smallest diameter tubes (6Fr) a smaller flush volume of 2mL may be used to support the use in very young children where fluid intake restriction may be of relevance.

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

Please consult the SmPC for full prescribing information and administration details.



ROS000019-009 May 2023

Rosemont Pharmaceuticals Ltd. Rosemont House, Yorkdale Industrial Park, Braithwaite Street, Leeds LS11 9XE **T** +44 (0)113 244 1400 **F** +44 (0)113 245 3567 **E** infodesk@rosemontpharma.com Sales/ Customer Service: **T** +44 (0) 113 244 1999 **F** +44 (0) 113 246 0738 **W** www.rosemontpharma.com

Adverse events should be reported. Reporting forms and information can be found at www.mhra.gov.uk/yellowcard Adverse events should also be reported to Rosemont Pharmaceuticals Ltd on 0113 244 1400.